



## The Duality Theorem for Min-Max Functions

Stéphane Gaubert\*, Jeremy Gunawardena  
Basic Research Institute in the Mathematical Sciences  
HP Laboratories Bristol  
HPL-BRIMS-97-16  
August, 1997

cycle time,  
generalized  
eigenvector,  
nonexpansive map,  
policy improvement

The set of min-max functions  $F: \mathbb{R}^n \rightarrow \mathbb{R}^n$  is the least set containing coordinate substitutions and translations and closed under pointwise max, min, and function composition. The Duality Conjecture asserts that the trajectories of a min-max function, considered as a dynamical system, have a linear growth rate (cycle time) and shows how this can be calculated through a representation of  $F$  as an infimum of max-plus linear functions. We prove the conjecture using an analogue of Howard's policy improvement scheme, carried out in a lattice ordered group of germs of affine functions at infinity. The methods yield an efficient algorithm for computing cycle times.

\*INRIA, Domaine de Voluceau, B.P. 105, 78153 Le Chesnay Cedex, France  
Internal Accession Date Only

# THE DUALITY THEOREM FOR MIN-MAX FUNCTIONS

STÉPHANE GAUBERT AND JEREMY GUNAWARDENA

**ABSTRACT.** The set of min-max functions  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is the least set containing coordinate substitutions and translations and closed under pointwise max, min, and function composition. The Duality Conjecture asserts that the trajectories of a min-max function, considered as a dynamical system, have a linear growth rate (cycle time) and shows how this can be calculated through a representation of  $F$  as an infimum of max-plus linear functions. We prove the conjecture using an analogue of Howard's policy improvement scheme, carried out in a lattice ordered group of germs of affine functions at infinity. The methods yield an efficient algorithm for computing cycle times.

## LE THÉORÈME DE DUALITÉ POUR LES FONCTIONS MIN-MAX

**RÉSUMÉ.** L'ensemble des fonctions min-max  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  est le plus petit ensemble de fonctions qui contient les substitutions de coordonnées et les translations, et qui est stable par les opérations min et max (point par point), ainsi que par composition. La Conjecture de Dualité affirme que les trajectoires d'un système récurrent gouverné par une dynamique min-max ont un taux de croissance linéaire (temps de cycle), qui se calcule à partir d'une représentation de  $F$  comme infimum de fonctions max-plus linéaires. La preuve utilise une itération sur les politiques à la Howard, à valeurs dans un groupe réticulé de germes de fonctions affines à l'infini. On a ainsi un algorithme efficace pour calculer le temps de cycle.

## 1. MIN-MAX FUNCTIONS

Let  $n$  denote a positive integer. We equip  $\mathbb{R}^n$  and functions  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  with the usual (pointwise) partial order. The least upper bound and greatest lower bound are denoted  $\vee$  and  $\wedge$ , respectively. We call *substitution* a function  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  of the form  $F(x)_i = x_{\pi(i)}$ , where  $\pi$  is an arbitrary (possibly non-bijective) transformation of  $\{1, \dots, n\}$ . A *translation* is a map  $\mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $x \mapsto x + u$ , with  $u \in \mathbb{R}^n$ . The set of min-max functions is the least set of functions  $\mathbb{R}^n \rightarrow \mathbb{R}^n$ , which contains substitutions and translations, and is stable by the operations  $\vee, \wedge$  and function composition.

Min-max functions include max-plus linear maps (i.e. linear maps over the max-plus semiring [6, 2, 19, 10]), which can be written as  $G(x)_i = \max_{1 \leq j \leq n} (A_{ij} + x_j)$ , where  $A$  is a  $n \times n$  matrix with entries in  $\mathbb{R} \cup \{-\infty\}$ , with at least one finite entry per row. Min-plus linear maps, defined dually, are also special min-max functions.

Using the mutual distributivity of  $\vee$  and  $\wedge$ , we can write any min-max function as

$$(1) \quad F(x) = \bigwedge_{G \in \mathcal{G}} G(x) = \bigvee_{H \in \mathcal{H}} H(x) ,$$

where  $\mathcal{G}$  and  $\mathcal{H}$  are finite sets of max-plus and min-plus linear maps, respectively.

Clearly, a min-max function  $F$  is *non-expansive* for the sup-norm  $\|\cdot\|$  (i.e.  $\|F(x) - F(y)\| \leq \|x - y\|$ ), it is *monotone* (i.e.  $x \leq y \implies F(x) \leq F(y)$ ), and (additively) *homogeneous* (i.e.  $F(x_1 + \alpha, \dots, x_n + \alpha) = F(x_1, \dots, x_n) + \alpha$ ).

Consider the trajectory  $\xi(k) = F(\xi(k-1))$  for  $k \geq 1$ ,  $\xi(0) = x \in \mathbb{R}^n$ . The *cycle-time* vector of  $F$ ,  $\chi(F)$ , is defined by

$$(2) \quad \chi(F) = \lim_{k \rightarrow \infty} \frac{1}{k} \times \xi(k) .$$

Since  $F$  is nonexpansive, the limit, if it exists, is independent of the initial point  $x$ . The existence of the cycle time vector for max-plus linear maps or dually, for min-plus linear maps, is stated in [12,

---

*Date:* June 20, 1997. Projet de note aux C.R.A.S.

Prop 2.1]. It is related with the max-plus Perron-Frobenius theory (see e.g. [2, §3.2.4,§3.7],[19],[10, § 3.7]).

A set  $\mathcal{S}$  of min-max functions is *rectangular* if for all  $G, G' \in \mathcal{S}$ , and for all  $i = 1, \dots, n$ , the function obtained by replacing the  $i$ -th component of  $G$  by the  $i$ -th component of  $G'$  belongs to  $\mathcal{S}$ . We denote by  $\overline{\mathcal{S}}$  the *rectangular closure* (least rectangular supset) of a set  $\mathcal{S}$ , which is finite when  $\mathcal{S}$  is finite. The min-max function (1) can be written equivalently as:

$$(3) \quad F(x) = \bigwedge_{G \in \overline{\mathcal{G}}} G(x) = \min_{G \in \overline{\mathcal{G}}} G(x) = \bigvee_{H \in \overline{\mathcal{H}}} H(x) = \max_{H \in \overline{\mathcal{H}}} H(x) ,$$

where “min” and “max” mean that the greatest lower bound and least upper bound are attained. Since min-max functions are monotone, we infer from (3) that

$$(4) \quad \bigwedge_{G \in \overline{\mathcal{G}}} \chi(G) \geq \bar{x} \geq \bigvee_{H \in \overline{\mathcal{H}}} \chi(H) ,$$

for any accumulation point  $\bar{x}$  of the sequence  $\frac{1}{k} \times \xi(k)$ . The *duality conjecture*, proposed by Gunawardena in [13], states that the extreme sides in (4) coincide. Hence,  $\chi(F) = \bar{x}$  exists, and its value is given explicitly by (4).

The Duality Conjecture was proved for  $n = 2$  in [4]. The cycle time vector was shown to exist for  $n = 3$  in [22]. The purpose of this note is to prove the duality conjecture, and to show how the cycle time can be computed efficiently.

Min-max functions were introduced in [14] following earlier work on special cases in [21]. Functions which are homogeneous and nonexpansive in the supremum norm —so called topical functions— have appeared in the work of several authors, in particular [5, 20, 16, 18]. They include (after suitable transformation) nonnegative matrices, Leontieff substitution systems, Bellman operators of games and of Markov decision processes with average cost and dynamical models of discrete event systems (digital circuits, computer networks, manufacturing plants, etc). Min-max functions play an important role within this larger class because any topical function  $T$  can be approximated by min-max functions in such a way that some of the dynamical behaviour of  $T$  is inherited by its approximations [17]. The cycle time vector appears in this context as a nonlinear generalisation of the conventional spectral radius of a nonnegative matrix. However, it does not always exist [16], and it is an important open problem to determine those topical functions for which it does. The positive solution of the Duality Conjecture shows that it does exist for min-max functions. In earlier work [4], we also showed that the Duality Conjecture implied the following strong fixed point theorem:  $F(x_1, \dots, x_n) = (x_1 + \alpha, \dots, x_n + \alpha)$  iff  $\chi(F) = (\alpha, \dots, \alpha)^T$ , which subsumes earlier results in the literature.

The lattice ordered group of germs  $\mathbb{G}$  introduced below is a variant of the semifield of asymptotic expansions introduced by Finkelstein and Roytberg, for its applications to statistical mechanics [8]. The same structure (semifield of max-jets) plays a key role in the study of asymptotics of the ordinary Perron eigenvalues and eigenvectors of matrices  $(A_{ij}^p)$ , where  $A_{ij} \geq 0$  and  $p \rightarrow \infty$  (see [1]). The algorithm below is an extension to germs of the policy iteration algorithm given in [4] (see also [3]), for the special min-max functions whose policies have an additive eigenvector. Classical policy iteration algorithms à la Howard (for stochastic control) can be found e.g. in [23].

## 2. GENERALIZED EIGENVECTORS AND GERMS OF AFFINE FUNCTIONS

We define an equivalence relation on  $\mathbb{R}^N$  by  $f \sim g \iff \exists K \in \mathbb{N}, \forall k \geq K, f(k) = g(k)$ . A map  $f \in \mathbb{R}^N$  is *affine* if  $f(k) = a + \alpha k$ , with  $a, \alpha \in \mathbb{R}$ . We denote by  $\mathbb{G}$  the image of the set of affine functions by the canonical projection:  $\mathbb{R}^N \rightarrow \mathbb{R}^N / \sim$ . An element of  $\mathbb{G}$  is a *germ of affine functions at  $+\infty$* . The usual laws  $\vee, \wedge, +, \times$  on  $\mathbb{R}^N$  induce quotient laws on  $\mathbb{R}^N / \sim$ . In particular,  $(\mathbb{G}, \vee, \wedge, +)$  is a lattice (totally) ordered group. We denote by  $\omega$  the equivalence class of the identity map  $\mathbb{N} \rightarrow \mathbb{R}, k \mapsto k$ . An element  $u$  of  $\mathbb{G}$  can be written in a unique way as  $u = a + \alpha\omega$ , with  $a, \alpha \in \mathbb{R}$ . We call  $\ell(u) \stackrel{\text{def}}{=} \alpha$  the *linear part* of  $u$ . E.g.  $((3 + 2\omega) \vee (2 + 5\omega)) \wedge (-100 + 7\omega) = 2 + 5\omega$ , for

$((3 + 2k) \vee (2 + 5k)) \wedge (-100 + 7k) = 2 + 5k$ , for  $k$  large enough. The shift operator  $\theta : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ ,  $\theta u(k) = u(k+1)$  induces a map  $\theta : \mathbb{G} \rightarrow \mathbb{G}$ ,  $\theta(a + \alpha\omega) = a + \alpha + \alpha\omega$ . The notation  $\theta, \ell, \vee, \wedge, +$  will be extended to vectors (entrywise). A min-max function  $F$  defines a map (also denoted  $F$ )  $\mathbb{G}^n \rightarrow \mathbb{G}^n$ , that to the equivalence class of  $u$ , associates the equivalence class of  $F \circ u$ .

A *generalized eigenvector* of  $F$  is a vector  $u \in \mathbb{G}^n$  such that  $F(u) = \theta u$ . More explicitly, if  $u = a + \alpha\omega$ , with  $a, \alpha \in \mathbb{R}^n$ ,

$$(5) \quad \exists K, \forall k \geq K, F(a + k\alpha) = a + (k+1)\alpha .$$

When  $\alpha = (\lambda, \dots, \lambda)^T$  with  $\lambda \in \mathbb{R}$ ,  $a$  is a (nonlinear, additive) eigenvector of  $F$ , with (nonlinear, additive) eigenvalue  $\lambda$ .

**THEOREM 1.** *Any min-max function has a generalized eigenvector.*

Before sketching the proof technique, we mention two immediate consequences.

**COROLLARY 1.** *Any min-max function has a cycle time.*

*Proof.* Let  $u = a + \alpha\omega$  denote a generalized eigenvector of a min-max function  $F$ . By (5), setting  $\xi(0) = a + K\alpha$ , we obtain  $\xi(k) = F(\xi(k-1)) = a + (k+K)\alpha$ . Hence,  $\chi(F) = \ell(u) = \alpha$ .  $\square$

**COROLLARY 2.** *The duality conjecture holds.*

*Proof.* Let  $u$  denote a generalized eigenvector of the min-max function  $F$  given by (1). Since  $\mathbb{G}$  is totally ordered and  $\overline{\mathcal{G}}$  is rectangular,  $\theta u = F(u) = G(u)$  for some  $G \in \overline{\mathcal{G}}$ . Hence,  $\chi(F) = \chi(G) = \ell(u)$ . This shows that the first relation in (4) is an equality. By symmetry, the second relation is also an equality, and the duality conjecture holds.  $\square$

The proof of Theorem 1, given in [9], relies on a policy iteration argument, which is analogous to Howard's multichain policy iteration algorithm for Markov decision processes with average cost (see e.g. [23]). Whereas the conventional policy iteration algorithm solves essentially a sequence of generalized spectral problems of the form  $c^{(k)} + P^{(k)}u = \theta u$ , with  $u \in \mathbb{G}^n$  (at step  $k$ ,  $c^{(k)} \in \mathbb{R}^n$  is given, and  $P^{(k)}$  is a given  $n \times n$  row-stochastic matrix), the min-max policy iteration algorithm below solves a sequence of generalized spectral problems which are linear over the max-plus semiring. The proof that it terminates is based on the following max-plus analogue of the maximum principle.

**LEMMA 1** (Max-plus Maximum Principle). *Let  $G$  denote a max-plus linear map  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $\chi(G) = 0$ . Let  $z \in \mathbb{R}^n$  such that  $G(z) \leq z$ . Let  $\mathcal{C}$  denote the set of nodes of the critical graph [2, §3.2.4, §3.7],[10, § 3.7] of  $G$ . There exists a unique  $x \in \mathbb{R}^n$  which is a solution of the “Dirichlet problem”*

$$\begin{aligned} G(x)_i &= x_i, \quad \forall i \notin \mathcal{C} , \\ x_i &= z_i, \quad \forall i \in \mathcal{C} . \end{aligned}$$

Furthermore,  $x \leq z$ .

If  $G$  is a max-plus linear map, we set  $\tilde{G} = -\chi(G) + G$ . The *spectral projector*  $P_G$  is defined by  $P_G(x) = \limsup_{k \rightarrow \infty} \tilde{G}^k(x)$ . When  $z$  and  $x$  are as in Lemma 1, we can check that  $x = P_G(z)$ . Hence, the max-plus maximum principle implies that  $u^{(k+1)} \leq u^{(k)}$  at each type (4b) step of Algorithm 1 below. The cycle time  $\chi(G)$ , the spectral projector, and a generalized eigenvector of  $G$ , can be computed in  $O(n^3)$  time (this follows essentially from [2, § 3.2.4, § 3.7]). Consider  $F$  as in (1). We will call *policies* the maps  $G \in \overline{\mathcal{G}}$ . The  $i$ -th component of a policy  $G$  is denoted  $G_i$ .

**ALGORITHM 1** (Min-max policy iteration). *Input:* A min-max function  $F$ . *Output:*  $u \in \mathbb{G}^n$  such that  $F(u) = \theta u$ .

1. *Initialization.* Select an arbitrary policy  $G^{(1)} \in \overline{\mathcal{G}}$ . Set  $k = 1$ . Find  $u^{(1)} \in \mathbb{G}^n$  such that  $G^{(1)}u^{(1)} = \theta u^{(1)}$ .

2. Let

$$\begin{aligned} I_k &= \{i \in \{1, \dots, n\} \mid F(u^{(k)})_i < (\theta u^{(k)})_i\} \\ J_k &= \{i \in \{1, \dots, n\} \mid \ell(F(u^{(k)}))_i < \ell(u^{(k)})_i\}. \end{aligned}$$

If  $I_k = \emptyset$ , we have  $F(u^{(k)}) = \theta u^{(k)}$ . Stop.

3. *Policy improvement.* We select  $G' \in \bar{\mathcal{G}}$  such that  $G'(u^{(k)}) = F(u^{(k)})$ .

(a) If  $J_k \neq \emptyset$ , we set:

$$G_i^{(k+1)} = \begin{cases} G'_i & \text{if } i \in J_k, \\ G_i^{(k)} & \text{if } i \notin J_k. \end{cases}$$

(b) If  $J_k = \emptyset$ , we set

$$G_i^{(k+1)} = \begin{cases} G'_i & \text{if } i \in I_k, \\ G_i^{(k)} & \text{if } i \notin I_k. \end{cases}$$

4. *Value determination.* Find  $v \in \mathbb{G}^n$  such that  $G^{(k+1)}(v) = \theta v$ .

(a) if  $\ell(v) \neq \ell(u^{(k)})$ , then, set  $u^{(k+1)} = v$ .

(b) if  $\ell(v) = \ell(u^{(k)})$ , then, select  $u^{(k+1)} = P_{G^{(k+1)}} u^{(k)}$ .

5. Increment  $k$  by one and go to step 2.

*Example 1.* We apply Algorithm 1 to:

$$F : \mathbb{R}^3 \rightarrow \mathbb{R}^3, F(x) = \begin{pmatrix} ((x_2 + 2) \vee (x_3 + 5)) \wedge x_1 \\ (x_2 + 1) \wedge (x_3 + 2) \\ (x_1 - 1) \vee (x_2 + 3) \end{pmatrix}.$$

We start from

$$G^{(1)}(x) = \begin{pmatrix} (x_2 + 2) \vee (x_3 + 5) \\ x_3 + 2 \\ (x_1 - 1) \vee (x_2 + 3) \end{pmatrix}, \quad u^{(1)} = \begin{pmatrix} 3 + 2.5\omega \\ 2.5\omega \\ 0.5 + 2.5\omega \end{pmatrix}.$$

This is a type (3b) policy improvement: we get  $F(u^{(1)}) = G^{(2)}(u^{(1)})$  with

$$G^{(2)}(x) = \begin{pmatrix} x_1 \\ x_2 + 1 \\ (x_1 - 1) \vee (x_2 + 3) \end{pmatrix}, \quad u^{(2)} = \begin{pmatrix} 0 \\ 1\omega \\ 2 + 1\omega \end{pmatrix}.$$

Then the algorithm terminates since  $F(u^{(2)}) = \theta u^{(2)}$ . Thus,  $\chi(F) = \ell(u^{(2)}) = (0, 1, 1)^T$ .

To conclude, we propose an extension of the duality conjecture. We will call *min-max-* $\mathbb{E}$  function a map of the form  $F(x) = \bigwedge_{i \in I} \bigvee_{j \in J} (c^{ij} + P^{ij}x)$ , where  $I, J$  are finite sets, and for all  $(i, j) \in I \times J$ :  $c^{ij} \in \mathbb{R}^n$  and  $P^{ij}$  is a probability matrix. When the  $P^{ij}$  matrices have exactly one nonzero entry per row, we obtain min-max functions.

**CONJECTURE 1.** *Any min-max-* $\mathbb{E}$  *function admits a generalized eigenvector.*

#### VERSION FRANÇAISE ABRÉGÉE

Soit  $n$  un entier positif. Nous munissons  $\mathbb{R}^n$  et l'ensemble des fonctions  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  de l'ordre partiel usuel (composante par composante). Les bornes sup et inf sont notées  $\vee$  et  $\wedge$ , respectivement. Nous appellerons *substitution* une application de la forme  $F(x)_i = x_{\pi(i)}$  où  $\pi$  est une transformation quelconque de  $\{1, \dots, n\}$  (peut-être non bijective). Une *translation* est une application de la forme:  $\mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $x \mapsto x + u$ , avec  $u \in \mathbb{R}^n$ . L'ensemble des fonctions min-max est le plus petit ensemble de fonctions  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  contenant les substitutions et les translations, et qui est stable par les opérations  $\vee, \wedge$  et la composition.

Les fonctions min-max comprennent les applications max-plus linéaires, (cf. [6, 2, 19, 10]), qui sont de la forme  $G(x)_i = \max_{1 \leq j \leq n} (A_{ij} + x_j)$ ,  $A$  étant une matrice  $n \times n$  à coefficients dans  $\mathbb{R} \cup \{-\infty\}$ , avec au moins un coefficient fini par ligne. Elles comprennent aussi les applications min-plus linéaires, définies dualement.

Comme  $\vee$  et  $\wedge$  distribuent mutuellement, nous pouvons écrire toute fonction min-max sous la forme (1), où  $\mathcal{G}$  et  $\mathcal{H}$  sont des ensembles finis d'applications max-plus et min-plus linéaires, respectivement.

On appelle vecteur de *temps de cycle* de  $F$  la quantité (2), avec  $\xi(k) = F(\xi(k-1))$  pour  $k \geq 1$  et  $\xi(0) = x$ . Quand elle existe, cette limite ne dépend pas du point initial  $x$ , car les fonctions min-max n'augmentent pas la distance associée à la norme-sup. Pour les applications max-plus ou min-plus linéaires, l'existence et le calcul du temps de cycle sont explicités dans [12]. Cela résulte de l'analogue max-plus de la théorie de Perron-Frobenius (voir par exemple [2, §3.2.4, §3.7],[19],[10, § 3.7]).

Un ensemble  $\mathcal{S}$  de fonctions min-max est *rectangulaire* si pour tout  $G, G' \in \mathcal{S}$ , et pour tout  $i = 1, \dots, n$ , la fonction obtenue en remplaçant la  $i$ -ème ligne de  $G$  par celle de  $G'$  appartient encore à  $\mathcal{S}$ . Nous noterons  $\overline{\mathcal{S}}$  la *clôture rectangulaire* (plus petit sur-ensemble rectangulaire) de  $\mathcal{S}$ . Évidemment, on peut prendre le sup ou l'inf sur la clôture rectangulaire sans changer la valeur de  $F$ , ce qui donne (3). Comme les fonctions min-max sont monotones, on déduit aisément (4) de (3), pour tout point d'accumulation  $\bar{x}$  de la suite  $\frac{1}{k} \times \xi(k)$ . La *conjecture de dualité*, énoncée par Gunawardena dans [13], affirme que les membres extrêmes de (4) sont égaux. Elle entraîne l'existence du temps de cycle  $\chi(F)$ .

Nous prouvons ici la conjecture de dualité, et montrons comment calculer efficacement le temps de cycle.

Motivations et références sur les fonctions min-max peuvent être trouvées dans [14, 21, 11, 4]. Le groupe réticulé des germes de fonctions affine  $\mathbb{G}$  employé ici est une variante du semicorps de développements asymptotiques introduit dans [8]. Voir aussi [1]. L'algorithme que nous donnons étend aux germes l'algorithme d'itération sur les politiques donné dans [4] (voir aussi [4]) pour les fonctions min-max particulières dont les politiques ont des vecteurs propres additifs.

Nous définissons la relation d'équivalence sur  $\mathbb{R}^N$ ,  $f \sim g \iff \exists K \in \mathbb{N}, \forall k \geq K, f(k) = g(k)$ . Nous notons  $\mathbb{G}$  l'ensemble des germes de fonctions affine en  $+\infty$ , i.e. l'image de l'ensemble des fonctions affines par la projection canonique:  $\mathbb{R}^N \rightarrow \mathbb{R}^N / \sim$ . Les lois usuelles  $\vee, \wedge, +, \times$  sur  $\mathbb{R}^N$  définissent des lois quotients sur  $\mathbb{R}^N / \sim$ . En particulier,  $(\mathbb{G}, \vee, \wedge, +)$  est un groupe totalement ordonné en treillis. Nous notons  $\omega$  la classe d'équivalence de l'application identité  $\mathbb{N} \rightarrow \mathbb{R}, k \mapsto k$ . Un germe  $u \in \mathbb{G}$  s'écrit de façon unique  $u = a + \alpha\omega$ , où  $a, \alpha \in \mathbb{R}$ , et  $\ell(u) \stackrel{\text{déf}}{=} \alpha$  est la *partie linéaire* de  $u$ . Ainsi,  $((3+2\omega) \vee (2+5\omega)) \wedge (-100+7\omega) = 2+5\omega$ , car  $((3+2k) \vee (2+5k)) \wedge (-100+7k) = 2+5k$ , quand  $k$  est assez grand. Le décalage  $\theta : \mathbb{R}^N \rightarrow \mathbb{R}^N$ ,  $\theta u(k) = u(k+1)$  induit une application  $\theta : \mathbb{G} \rightarrow \mathbb{G}$ ,  $\theta(a + \alpha\omega) = a + \alpha + \alpha\omega$ . Nous entendrons les notations  $\theta, \ell, \vee, \wedge, +$  aux vecteurs, composante par composante. Une fonction min-max  $F$  définit une application  $\mathbb{G}^n \rightarrow \mathbb{G}^n$ ,  $u \mapsto F \circ u$ , que nous noterons encore  $F$ .

Un *vecteur propre généralisé* de  $F$  est un vecteur  $u \in \mathbb{G}^n$  tel que  $F(u) = \theta u$ . Si  $u = a + \alpha\omega$ , avec  $a, \alpha \in \mathbb{R}^n$ , cela s'écrit comme en (5).

**THÉORÈME 1.** *Toute fonction min-max admet un vecteur propre généralisé.*

**COROLLAIRE 1.** *Toute fonction min-max admet un temps de cycle.*

**COROLLAIRE 2.** *La conjecture de dualité est satisfaite.*

La preuve du théorème 1, donnée dans [9], repose sur un argument d'amélioration des politiques à la Howard. La preuve de convergence utilise un principe du maximum max-plus. Nous donnons seulement ici l'algorithme qui résulte de cette preuve. Pour une application max-plus linéaire  $G$ , on note  $\tilde{G} = -\chi(G) + G$ . Lorsque  $\chi(G) = 0$ , le *projecteur spectral*  $P_G$  de  $G$  est défini par  $P_G(x) = \limsup_{k \rightarrow \infty} \tilde{G}^k(x)$ . Signalons que les divers objets spectraux d'une application max-plus linéaire peuvent être calculés en temps  $\mathcal{O}(n^3)$  (voir [2, §3.2.4, §3.7]). Soit  $F$  comme en (1). Nous appellerons *politiques* les applications  $G \in \overline{\mathcal{G}}$ . L'algorithme 1 procède comme suit. On part d'une politique arbitraire  $G^{(1)} \in \mathcal{G}$  et d'un vecteur propre généralisé associé  $u^{(1)}$ . Si  $F(u^{(1)}) = \theta u^{(1)}$ , on s'arrête. Sinon, on choisit  $G' \in \overline{\mathcal{G}}$  tel que  $G'(u^{(1)}) = F(u^{(1)})$ , et l'on construit une nouvelle politique  $G^{(2)}$  selon les cas (3a) et (3b), On calcule un vecteur propre généralisé  $u^{(2)}$  de  $G^{(2)}$  (dans le cas (4b), on doit choisir  $u^{(2)} = P_{G^{(2)}} u^{(2)}$ ). Et ainsi de suite.

Concluons par une extension de la conjecture de dualité. Nous appellerons *fonction min-max- $\mathbb{E}$*  toute application de la forme  $F(x) = \bigwedge_{i \in I} \bigvee_{j \in J} (c^{ij} + P^{ij}x)$ , où  $I, J$  sont des ensembles finis, avec pour tout  $(i, j) \in I \times J$ ,  $c^{ij} \in \mathbb{R}^n$ , et  $P^{ij}$  matrice de probabilité. Quand les matrices  $P^{ij}$  ont exactement un coefficient non-nul par ligne, on obtient les fonctions min-max.

**CONJECTURE 1.** *Toute fonction min-max- $\mathbb{E}$  admet un vecteur propre généralisé.*

## REFERENCES

- [1] M. Akian, R.B. Bapat, and S. Gaubert. Asymptotics of the Perron eigenvalue and eigenvector using max algebra. In preparation, 1997.
- [2] F. Baccelli, G. Cohen, G.J. Olsder, and J.P. Quadrat. *Synchronization and Linearity*. Wiley, 1992.
- [3] J. Cochet-Terrasson and S. Gaubert. A min-max analogue of the Howard algorithm. Privately circulated draft. Appears in the appendix of: J. Cochet-Terrasson, Rapport de Stage de DEA "Modélisation et Méthodes Mathématiques en Économie", Université de Paris I, July 1996.
- [4] J. Cochet-Terrasson, S. Gaubert, and J. Gunawardena. Dynamics of min-max functions. Submitted for publication, 1997.
- [5] M.G. Crandall and L. Tartar. Some relations between non expansive and order preserving maps. *Proceedings of the AMS*, 78(3):385–390, 1980.
- [6] R.A. Cuninghame-Green. *Minimax Algebra*. Number 166 in Lecture notes in Economics and Mathematical Systems. Springer, 1979.
- [7] R.A Cuninghame-Green. Minimax algebra and applications. *Advances in Imaging and Electron Physics*, 90, 1995.
- [8] A.V. Finkelstein and M.A. Roitberg. Computation of biopolymers: a general approach to different problems. *BioSystems*, 30:1–20, 1993.
- [9] S. Gaubert and J. Gunawardena. A proof of the duality conjecture for min-max functions. Manuscrit joint à la présente note aux C.R.A.S., 1997.
- [10] S. Gaubert and M. Plus. Methods and applications of  $(\max, +)$  linear algebra. In R. Reischuk and M. Morvan, editors, *STACS'97*, number 1200 in LNCS, Lübeck, March 1997. Springer.
- [11] J. Gunawardena. Timing analysis of digital circuits and the theory of min-max functions. In *Digest of Technical Papers of the ACM International Workshop on Timing Issues in the Specification and Synthesis of Digital Systems*. ACM, 1993.
- [12] J. Gunawardena. Cycle time and fixed points of min-max functions. In G. Cohen and J.P. Quadrat, editors, *11th International Conference on Analysis and Optimization of Systems*, number 199 in LNCIS, pages 266–272. Springer, 1994.
- [13] J. Gunawardena. Cycle time and fixed points of min-max functions. In G. Cohen and J.P. Quadrat, editors, *11th International Conference on Analysis and Optimization of Systems*, number 199 in LNCIS, pages 266–272. Springer, 1994.
- [14] J. Gunawardena. Min-max functions. *J. Discrete Event Dynamic Systems*, 4:377–406, 1994.
- [15] J. Gunawardena, editor. *Idempotency*. Publications of the Newton Institute. Cambridge University Press, 1997.
- [16] J. Gunawardena and M. Keane. On the existence of cycle times for some nonexpansive maps. Technical Report HPL-BRIMS-95-003, Hewlett-Packard Labs, 1995.
- [17] J. Gunawardena, M. Keane, and C. Sparrow. In preparation, 1997.
- [18] V. Kolokoltsov. Linear additive and homogenears operators. Appears in [19].
- [19] V. Maslov and S. Samborskii, editors. *Idempotent analysis*, volume 13 of *Adv. in Sov. Math.* AMS, RI, 1992.
- [20] R.D. Nussbaum. Hilbert's projective metric and iterated nonlinear maps. *Memoirs of the AMS*, 95(391), 1990.
- [21] G.J. Olsder. Eigenvalues of dynamic max-min systems. *J. Discrete Event Dynamic Systems*, 1:177–207, 1991.
- [22] C. Sparrow. Existence of cycle time for minmax functions of dimension 3. Technical Report HPL-BRIMS-96-008, Hewlett-Packard Labs, 1996.
- [23] P. Whittle. *Optimization over Time*. Wiley, 1986.

INRIA, DOMAINE DE VOLUCEAU, B.P. 105, 78153 LE CHESNAY CEDEX, FRANCE. PHONE. +33 1 39 63 52 58; FAX. +33 1 39 63 57 86

*E-mail address:* Stephane.Gaubert@inria.fr

BRIMS, HEWLETT-PACKARD LABS, FILTON ROAD, STOKE GIFFORD, BRISTOL BS12 6QZ, UK

*E-mail address:* jhcg@hplb.hpl.hp.com